

An algebraic formula for the intersection number of a polynomial immersion

Iwona Karolkiewicz Aleksandra Nowel
Zbigniew Szafraniec

July 2007

Abstract

There is presented an algorithm for computing the topological degree for a large class of polynomial mappings. As an application there is given an effective algebraic formula for the intersection number of a polynomial immersion $M \rightarrow \mathbb{R}^{2m}$, where M is an m -dimensional algebraic manifold.

Keywords: polynomial immersion, intersection number, local topological degree

Let $H = (h_1, \dots, h_n) : \mathbb{R}^n \longrightarrow \mathbb{R}^n$ be a continuous mapping, and let p be isolated in $H^{-1}(0)$. Then one may define the local topological degree $\deg_p H$ as the topological degree of the mapping

$$S_r^{n-1} \ni x \longmapsto H(x)/\|H(x)\| \in S^{n-1},$$

where S_r^{n-1} is the sphere of radius $r \ll 1$ centered at p .

Suppose that H is a polynomial mapping and $u : \mathbb{R}^n \longrightarrow \mathbb{R}$ is a polynomial. Put $U = \{x \in \mathbb{R}^n : u(x) > 0\}$. Let $J_{\mathbb{R}} \subset \mathbb{R}[x] = \mathbb{R}[x_1, \dots, x_n]$ denote the ideal generated by $h_1, \dots, h_n$, and let $Q = \mathbb{R}[x]/J_{\mathbb{R}}$. Suppose that $\dim_{\mathbb{R}} Q < \infty$, so that $H^{-1}(0)$ is finite. According to [10], one may construct quadratic forms Φ_T and Ψ_T on Q such that

$$\sum \deg_p H = (\text{signature } \Phi_T + \text{signature } \Psi_T)/2,$$

Iwona Karolkiewicz, Aleksandra Nowel and Zbigniew Szafraniec
University of Gdańsk, Institute of Mathematics
80-952 Gdańsk, Wita Stwosza 57
Poland
Tel.: +48-58-5232059
Fax: +48-58-3414914
Email: ikarolki@manta.univ.gda.pl
Email: Aleksandra.Nowel@math.univ.gda.pl
Email: Zbigniew.Szafraniec@math.univ.gda.pl

where $p \in H^{-1}(0) \cap U$. In [10] is also presented a simpler formula for

$$\sum \deg_p H \pmod{2}.$$

The assumption $\dim_{\mathbb{R}} Q < \infty$ is too restrictive in some cases. For instance if $H^{-1}(0)$ is infinite then $\dim_{\mathbb{R}} Q = \infty$.

This paper is devoted to the case where there exists an ideal $I_{\mathbb{R}} \supset J_{\mathbb{R}}$ such that

$$(J_{\mathbb{R}} : I_{\mathbb{R}}) + I_{\mathbb{R}} = \mathbb{R}[x] \quad \text{and} \quad \dim_{\mathbb{R}} \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}}) < \infty.$$

(The quotient of ideals $J_{\mathbb{R}} : I_{\mathbb{R}}$ is defined as the set of all $f \in \mathbb{R}[x]$ with $fg \in J_{\mathbb{R}}$ for all $g \in I_{\mathbb{R}}$.)

We shall show that in that case $H^{-1}(0) \setminus V(I_{\mathbb{R}}) = V(J_{\mathbb{R}} : I_{\mathbb{R}})$ is finite, and there are effective algebraic formulae for $\sum \deg_p H \pmod{2}$ and $\sum \deg_p H$ analogous to the ones proved in [10].

Let M be a compact oriented m -dimensional manifold ($m > 1$), and let $g : M \rightarrow \mathbb{R}^{2m}$ be an immersion. Whitney in [12] introduced the intersection number $I(g)$.

Assume that $M \subset \mathbb{R}^{n+m}$ is an algebraic complete intersection, and a polynomial immersion g has a finite set of self-intersections. Let $\Delta \subset \mathbb{R}^{n+m} \times \mathbb{R}^{n+m}$ be the diagonal.

We show that one may construct a polynomial mapping $H : \mathbb{R}^{n+m} \times \mathbb{R}^{n+m} \rightarrow \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{2m}$ such that $\Delta \subset H^{-1}(0)$, $H^{-1}(0) \setminus \Delta$ is finite, and for even m

$$I(g) = \frac{1}{2} \sum_p \deg_p H,$$

where $p \in H^{-1}(0) \setminus \Delta$. There is also presented a similar formula for odd $m > 1$.

As $\Delta \subset H^{-1}(0)$ is infinite, we cannot apply method presented in [10] so as to compute $\sum_p \deg_p H$. We show how to apply methods developed in this paper so as to express the intersection number $I(g)$ either as the signature of a quadratic form on $\mathcal{A} = \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}})$ (for even m), in terms of $\dim_{\mathbb{R}} \mathcal{A}$ and signs of determinants of matrices of two quadratic forms (for odd $m > 1$).

The paper is organized as follows. In Section 1 we show that the complex algebra $\mathbb{C}[x]/(J_{\mathbb{C}} : I_{\mathbb{C}})$ is isomorphic to the product of some algebras associated to the ideal $J_{\mathbb{C}}$ and points in $H_{\mathbb{C}}^{-1}(0) \setminus V(I_{\mathbb{C}})$. In Section 2 we discuss relations between the ideal $J_{\mathbb{R}} : I_{\mathbb{R}}$ and its complex counterpart $J_{\mathbb{C}} : I_{\mathbb{C}}$. In Section 3 we recall some results presented in [10]. In particular we explain how to construct the bilinear forms which we use in Section 4, where we prove a formula for $\sum_p \deg_p H$ in terms of their signatures. In Section 5 we apply these results so as to give an effective method for computing the intersection number of a polynomial immersion on an algebraic manifold. In the end we present examples computed by a computer. We implemented our algorithm with help of SINGULAR [4]. We have also used computer programs written by Adriana Gorzelak and Magdalena Sarnowska — students of computer sciences at the University of Gdańsk — which can construct matrices of Φ_T , Ψ_T , and compute the signature of a bilinear form.

1 Quotients of ideals in $\mathbb{C}[x]$

If S is an ideal in $\mathbb{C}[x] = \mathbb{C}[x_1, \dots, x_n]$ then by $V(S)$ we will denote set of all complex zeros of S . For each $p = (p_1, \dots, p_n) \in \mathbb{C}^n$, m_p is the maximal ideal generated by monomials $x_1 - p_1, \dots, x_n - p_n$. If $J, I \subset \mathbb{C}[x]$ are ideals, then

$$J : I = \{f \in \mathbb{C}[x] : fg \in J \text{ for all } g \in I\}$$

is called the ideal quotient of J by I . Since $J \subset J : I$, $V(J : I) \subset V(J)$.

Suppose that $J, I \subset \mathbb{C}[x]$ are ideals such that $J \subset I$ and

$$\dim_{\mathbb{C}} \frac{\mathbb{C}[x]}{J : I} < \infty.$$

Then $V(J : I)$ is finite. Applying [2, Theorem 7, p. 192] we get inclusions:

$$V(J) \setminus V(I) \subset \overline{V(J) \setminus V(I)} \subset V(J : I) \subset V(J),$$

so that $V(J) \setminus V(I) \subset V(J : I) \setminus V(I) \subset V(J) \setminus V(I)$. Denote $V(J) \setminus V(I) = V(J : I) \setminus V(I) = \{p_1, \dots, p_s\}$.

Proposition 1.1 *There exists a positive integer k such that*

$$\frac{\mathbb{C}[x]}{J} \cong \frac{\mathbb{C}[x]}{J + I^k} \times \frac{\mathbb{C}[x]}{J + m_{p_1}^k} \times \dots \times \frac{\mathbb{C}[x]}{J + m_{p_s}^k}.$$

In particular $J = (J + I^k) \cap (J + m_{p_1}^k) \cap \dots \cap (J + m_{p_s}^k)$. The same holds true for any integer bigger than k .

Proof. We have $V(I) \subset V(J)$ and $V(J) \setminus V(I) = \{p_1, \dots, p_s\}$, so

$$V(J) = \{p_1, \dots, p_s\} \cup V(I) = V(m_{p_1}) \cup \dots \cup V(m_{p_s}) \cup V(I) = V(m_{p_1} \cap \dots \cap m_{p_s} \cap I).$$

By the Hilbert Nullstellensatz, there exists k such that

$$(1.1) \quad (m_{p_1} \cap \dots \cap m_{p_s} \cap I)^k \subset J.$$

If $i \neq j$ then $V(m_{p_i}^k + m_{p_j}^k) = V(m_{p_i}^k) \cap V(m_{p_j}^k) = \emptyset$, so,

$$\mathbb{C}[x] = m_{p_i}^k + m_{p_j}^k \subseteq (J + m_{p_i}^k) + (J + m_{p_j}^k) \subseteq \mathbb{C}[x].$$

and then

$$(J + m_{p_i}^k) + (J + m_{p_j}^k) = \mathbb{C}[x].$$

We have $V(I^k + m_{p_i}^k) = V(I) \cap \{p_i\} = \emptyset$, so that

$$\mathbb{C}[x] = I^k + m_{p_i}^k \subseteq (J + I^k) + (J + m_{p_i}^k) \subseteq \mathbb{C}[x],$$

and then

$$(J + I^k) + (J + m_{p_i}^k) = \mathbb{C}[x].$$

The Chinese Remainder Theorem implies that:

$$\frac{\mathbb{C}[x]}{(J + I^k) \cap (J + m_{p_1}^k) \cap \dots \cap (J + m_{p_s}^k)} \cong \frac{\mathbb{C}[x]}{J + I^k} \times \frac{\mathbb{C}[x]}{J + m_{p_1}^k} \times \dots \times \frac{\mathbb{C}[x]}{J + m_{p_s}^k}$$

and

$$(J + I^k) \cap (J + m_{p_1}^k) \cap \dots \cap (J + m_{p_s}^k) = (J + I^k) \cdot (J + m_{p_1}^k) \cdot \dots \cdot (J + m_{p_s}^k).$$

By (1.1) we get

$$\begin{aligned} J &\subset (J + I^k) \cap (J + m_{p_1}^k) \cap \dots \cap (J + m_{p_s}^k) = \\ &= (J + I^k) \cdot (J + m_{p_1}^k) \cdot \dots \cdot (J + m_{p_s}^k) \subset J + (I^k \cdot m_{p_1}^k \cdot \dots \cdot m_{p_s}^k) = \\ &= J + (I \cdot m_{p_1} \cdot \dots \cdot m_{p_s})^k \subset J + (I \cap m_{p_1} \cap \dots \cap m_{p_s})^k \subset J + J, \end{aligned}$$

and then $(J + I^k) \cap (J + m_{p_1}^k) \cap \dots \cap (J + m_{p_s}^k) = J$. So we have

$$\frac{\mathbb{C}[x]}{J} \cong \frac{\mathbb{C}[x]}{J + I^k} \times \frac{\mathbb{C}[x]}{J + m_{p_1}^k} \times \dots \times \frac{\mathbb{C}[x]}{J + m_{p_s}^k}.$$

□

Lemma 1.2 *Let $p \in V(J)$. For any positive integer k , f is invertible in $\mathbb{C}[x]/(J + m_p^k)$ if and only if $f(p) \neq 0$.*

Proof. ($\Rightarrow$) Since f is invertible in $\mathbb{C}[x]/(J + m_p^k)$, there exists $g \in \mathbb{C}[x]$ such that $fg \equiv 1 \pmod{J + m_p^k}$, so

$$fg = 1 + w + v,$$

where $w \in J$ and $v \in m_p^k$. Hence

$$f(p)g(p) = 1 + w(p) + v(p) = 1,$$

and so $f(p) \neq 0$.

($\Leftarrow$) f can be presented as $f = f(p) - g$, where $g \in m_p$. Then

$$(f(p) - g)(f(p)^{k-1} + f(p)^{k-2}g + f(p)^{k-3}g^2 + \dots + g^{k-1}) = f(p)^k - g^k.$$

Set $h = (f(p)^{k-1} + f(p)^{k-2}g + \dots + g^{k-1})$. We have $g^k \in m_p^k$, so

$fh = (f(p) - g)h \equiv f(p)^k \neq 0$ in $\mathbb{C}[x]/(J + m_p^k)$, and then f is invertible in $\mathbb{C}[x]/(J + m_p^k)$. □

Take k as in Proposition 1.1. Then

Lemma 1.3 $J : I = (\bigcap_{i=1}^s (J + m_{p_i}^k)) \cap ((J + I^k) : I)$

Proof. Take $f \in \bigcap_{i=1}^s (J + m_{p_i}^k) \cap ((J + I^k) : I)$. For any $g \in I$ and $1 \leq i \leq s$ we have

$$fg \in J + I^k \quad \text{and} \quad fg \in J + m_{p_i}^k,$$

so $fg \in (J + I^k) \cap (J + m_{p_1}^k) \cap \dots \cap (J + m_{p_s}^k)$. Using Proposition 1.1 we get $fg \in J$, so that $f \in J : I$.

Now let us take $f \in J : I$. For any $g \in I$ we have $fg \in J \subset (J + I^k)$, and then $f \in (J + I^k) : I$. For $1 \leq i \leq s$ we also have

$$fg \in J + m_{p_i}^k.$$

Because $p_i \notin V(I)$, then there exists $h \in I$ such that $h(p_i) \neq 0$. From Lemma 1.2, h is invertible in $\mathbb{C}[x]/(J + m_{p_i}^k)$. As $fh \equiv 0$ in $\mathbb{C}[x]/(J + m_{p_i}^k)$ we have

$$f \in J + m_{p_i}^k.$$

So $f \in (\bigcap_{i=1}^s (J + m_{p_i}^k)) \cap ((J + I^k) : I)$. □

From now on we shall assume that $(J : I) + I = \mathbb{C}[x]$. We have

Lemma 1.4 *If $(J : I) + I = \mathbb{C}[x]$ and $\dim_{\mathbb{C}} \mathbb{C}[x]/(J : I) < \infty$ then $V(J : I) \cap V(I) = \emptyset$, and then $V(J : I) = V(J) \setminus V(I) = \{p_1, \dots, p_s\}$.*

□

Lemma 1.5 *For every positive integer k we have $(J : I) + I^k = \mathbb{C}[x]$.*

Proof. We have

$$V((J : I) + I^k) = V(J : I) \cap V(I^k) = V(J : I) \cap V(I).$$

Since $(J : I) + I = \mathbb{C}[x]$, we have $V(J : I) \cap V(I) = \emptyset$. So

$$(J : I) + I^k = \mathbb{C}[x].$$

□

Lemma 1.6 *For every positive integer k we have $(J + I^k) : I = \mathbb{C}[x]$.*

Proof. Let us take $f \in (J : I) + I^k$. Then $f = h_1 + h_2$, where $h_1 \in J : I$ and $h_2 \in I^k$. For any $g \in I$ we get

$$fg = h_1g + h_2g \in J + I^k.$$

So we have $(J : I) + I^k \subseteq (J + I^k) : I$. Using previous Lemma we get

$$\mathbb{C}[x] = (J : I) + I^k \subseteq (J + I^k) : I \subseteq \mathbb{C}[x].$$

□

Lemmas 1.3, 1.6 and Proposition 1.1 imply

Corollary 1.7 *If $(J : I) + I = \mathbb{C}[x]$ then for all k large enough*

$$J : I = \bigcap_{i=1}^s (J + m_{p_i}^k).$$

□

Ideals $J + m_{p_i}^k$ are pairwise comaximal. As a consequence of the Chinese Remainder Theorem we get

Proposition 1.8 *If $(J : I) + I = \mathbb{C}[x]$ then for all k large enough*

$$\frac{\mathbb{C}[x]}{J : I} \cong \frac{\mathbb{C}[x]}{J + m_{p_1}^k} \times \dots \times \frac{\mathbb{C}[x]}{J + m_{p_s}^k}.$$

□

2 Quotients of ideals in $\mathbb{R}[x]$ and $\mathbb{C}[x]$

Lemma 2.1 *Let $f_1, \dots, f_r$ be polynomials with real coefficients. Let $S_{\mathbb{R}}$ (resp. $S_{\mathbb{C}}$) denote the ideal in $\mathbb{R}[x]$ (resp. $\mathbb{C}[x]$) generated by $f_1, \dots, f_r$. Then*

- (i) $S_{\mathbb{R}} = S_{\mathbb{C}} \cap \mathbb{R}[x]$,
- (ii) $S_{\mathbb{R}} = \mathbb{R}[x] \Leftrightarrow S_{\mathbb{C}} = \mathbb{C}[x]$.

Proof.

- (i) If $f \in S_{\mathbb{R}}$ then obviously $f \in S_{\mathbb{C}} \cap \mathbb{R}[x]$.

If $h = \sum_{\alpha} a_{\alpha} x^{\alpha} \in \mathbb{C}[x]$ then put $\bar{h} = \sum_{\alpha} \overline{a_{\alpha}} x^{\alpha}$. In that case $h \in \mathbb{R}[x]$ if and only if $h = \bar{h}$. Take $f \in S_{\mathbb{C}} \cap \mathbb{R}[x]$. There exist $h_1, \dots, h_r \in \mathbb{C}[x]$ such that $f = \sum_{i=1}^r h_i f_i$. As $f, f_1, \dots, f_r \in \mathbb{R}[x]$, we have $\bar{f} = f = \sum_{i=1}^r \bar{h}_i f_i$. Hence $f = \frac{1}{2} \sum_{i=1}^r (h_i + \bar{h}_i) f_i$. Of course $h_i + \bar{h}_i \in \mathbb{R}[x]$, so $f \in S_{\mathbb{R}}$.

- (ii) If $S_{\mathbb{R}} = \mathbb{R}[x]$ then $1 \in S_{\mathbb{R}} \subset S_{\mathbb{C}}$, so $S_{\mathbb{C}} = \mathbb{C}[x]$.

If $S_{\mathbb{C}} = \mathbb{C}[x]$ then $S_{\mathbb{R}} = S_{\mathbb{C}} \cap \mathbb{R}[x] = \mathbb{C}[x] \cap \mathbb{R}[x] = \mathbb{R}[x]$.

□

Lemma 2.2 *Assume that $J_{\mathbb{R}}, I_{\mathbb{R}}$ are ideals in $\mathbb{R}[x]$. Then $(J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}} = J_{\mathbb{C}} : I_{\mathbb{C}}$, and then $(J_{\mathbb{C}} : I_{\mathbb{C}}) \cap \mathbb{R}[x] = J_{\mathbb{R}} : I_{\mathbb{R}}$.*

Proof. Let $g_1, \dots, g_s \in \mathbb{R}[x]$ be generators of the ideal $I_{\mathbb{R}}$, and in consequence of the ideal $I_{\mathbb{C}}$, and let $f_1, \dots, f_l$ generate $J_{\mathbb{R}}$ and also $J_{\mathbb{C}}$. Of course

$$J_{\mathbb{R}} : I_{\mathbb{R}} = \{h \in \mathbb{R}[x] : hg_i \in J_{\mathbb{R}}, \text{ for each } 1 \leq i \leq s\}$$

$$J_{\mathbb{C}} : I_{\mathbb{C}} = \{h \in \mathbb{C}[x] : hg_i \in J_{\mathbb{C}}, \text{ for each } 1 \leq i \leq s\}.$$

Take $h \in (J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}}$, then there exist $w_1, \dots, w_m \in \mathbb{C}[x]$ and $v_1, \dots, v_m \in J_{\mathbb{R}} : I_{\mathbb{R}}$ such that

$$h = \sum_{j=1}^m w_j v_j.$$

For each $1 \leq i \leq s$, $v_j g_i \in J_{\mathbb{R}}$, and $hg_i \in J_{\mathbb{C}}$, so $h \in J_{\mathbb{C}} : I_{\mathbb{C}}$.

Take $h \in J_{\mathbb{C}} : I_{\mathbb{C}}$, then for each $1 \leq i \leq s$, $hg_i \in J_{\mathbb{C}}$, so there exist $w_1, \dots, w_l \in \mathbb{C}[x]$ such that $hg_i = \sum_{j=1}^l f_j w_j$. Then $\bar{h}g_i = \sum_{j=1}^l f_j \bar{w}_j$ and

$$(h + \bar{h})g_i = \sum_{j=1}^l (w_j + \bar{w}_j) f_j.$$

Of course $w_j + \bar{w}_j \in \mathbb{R}[x]$. So $(h + \bar{h})g_i \in J_{\mathbb{R}}$ for each $1 \leq i \leq s$, and then $h + \bar{h} \in J_{\mathbb{R}} : I_{\mathbb{R}} \subset (J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}}$. As

$$(h - \bar{h})g_i = \sum_{j=1}^l (w_j - \bar{w}_j) f_j$$

and $\sqrt{-1}(w_j - \bar{w}_j) \in \mathbb{R}[x]$ so $\sqrt{-1}(h - \bar{h})g_i \in J_{\mathbb{R}}$ for each $1 \leq i \leq s$. So we get $\sqrt{-1}(h - \bar{h}) \in J_{\mathbb{R}} : I_{\mathbb{R}}$, and then $h - \bar{h} \in (J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}}$. Because $h + \bar{h} \in (J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}}$ and $h - \bar{h} \in (J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}}$ then $h \in (J_{\mathbb{R}} : I_{\mathbb{R}})_{\mathbb{C}}$. $\square$

If $S_{\mathbb{K}}$ is an ideal in $\mathbb{K}[x]$, where $\mathbb{K}$ is either $\mathbb{C}$ or $\mathbb{R}$, denote

$$V(S_{\mathbb{K}}) = \{p \in \mathbb{K}^n : f(p) = 0 \text{ for all } f \in S_{\mathbb{K}}\}.$$

Consider ideals $J_{\mathbb{R}} \subset I_{\mathbb{R}} \subset \mathbb{R}[x]$, such that

$$\dim_{\mathbb{R}} \frac{\mathbb{R}[x]}{J_{\mathbb{R}} : I_{\mathbb{R}}} < \infty \quad \text{and} \quad (J_{\mathbb{R}} : I_{\mathbb{R}}) + I_{\mathbb{R}} = \mathbb{R}[x].$$

Then $\mathbb{C}[x] = (J_{\mathbb{C}} : I_{\mathbb{C}}) + I_{\mathbb{C}}$ and $\dim_{\mathbb{C}} \mathbb{C}[x]/(J_{\mathbb{C}} : I_{\mathbb{C}}) = \dim_{\mathbb{R}} \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}}) < \infty$. By Lemma 1.4, $V(J_{\mathbb{C}} : I_{\mathbb{C}}) = V(J_{\mathbb{C}}) \setminus V(I_{\mathbb{C}})$ is finite. Let $V(J_{\mathbb{C}}) \setminus V(I_{\mathbb{C}}) = \{p_1, \dots, p_s\}$. By Proposition 1.8 there exists a positive integer k such that

$$\frac{\mathbb{C}[x]}{(J_{\mathbb{C}} : I_{\mathbb{C}})} \cong \frac{\mathbb{C}[x]}{J_{\mathbb{C}} + m_{p_1}^k} \times \dots \times \frac{\mathbb{C}[x]}{J_{\mathbb{C}} + m_{p_s}^k}.$$

Set $V(J_{\mathbb{R}}) \setminus V(I_{\mathbb{R}}) = \{p_1, \dots, p_m\}$ and $(V(J_{\mathbb{C}}) \setminus V(I_{\mathbb{C}})) \setminus \mathbb{R}^n = \{q_1, \bar{q}_1, \dots, q_r, \bar{q}_r\}$. Of course $s = m + 2r$. For k large enough and $p \in V(J_{\mathbb{R}}) \setminus V(I_{\mathbb{R}})$ we define an $\mathbb{R}$ -algebra

$$\mathcal{A}_{\mathbb{R},p} := \frac{\mathbb{R}[x]}{J_{\mathbb{R}} + m_{\mathbb{R},p}^k}, \quad \text{where } m_{\mathbb{R},p} = \{f \in \mathbb{R}[x] : f(p) = 0\}.$$

For $p \in V(J_{\mathbb{C}}) \setminus V(I_{\mathbb{C}})$, we define an $\mathbb{C}$ -algebra

$$\mathcal{A}_{\mathbb{C},p} := \frac{\mathbb{C}[x]}{J_{\mathbb{C}} + m_{\mathbb{C},p}^k}, \quad \text{where } m_{\mathbb{C},p} = \{f \in \mathbb{C}[x] : f(p) = 0\}.$$

Of course, $f \in J_{\mathbb{C}} + m_{\mathbb{C},p}^k$ if and only if $\bar{f} \in J_{\mathbb{C}} + m_{\mathbb{C},\bar{p}}^k$. In particular

$$(2.1) \quad \mathbb{R}[x] \cap (J_{\mathbb{C}} + m_{\mathbb{C},p}^k) = \mathbb{R}[x] \cap J_{\mathbb{C}} + m_{\mathbb{C},\bar{p}}^k.$$

The mapping $f \mapsto \bar{f}$ induces an $\mathbb{R}$ -isomorphism of algebras $\mathcal{A}_{\mathbb{C},p}$ and $\mathcal{A}_{\mathbb{C},\bar{p}}$, so that

$$(2.2) \quad \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C},p} = 2 \dim_{\mathbb{R}} \mathcal{A}_{\mathbb{C},p} = 2 \dim_{\mathbb{R}} \mathcal{A}_{\mathbb{C},\bar{p}} = \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C},\bar{p}}.$$

Let us denote

$$\mathcal{B} = \oplus_{i=1}^m \mathcal{A}_{\mathbb{R},p_i} \oplus_{j=1}^r \mathcal{A}_{\mathbb{C},q_j}.$$

Theorem 2.3 *For all k large enough there is a natural isomorphism*

$$\frac{\mathbb{R}[x]}{J_{\mathbb{R}} : I_{\mathbb{R}}} \cong \mathcal{B}.$$

Proof. We take k as in Proposition 1.8. We define a homomorphism

$$\pi : \mathbb{R}[x] \longrightarrow \mathcal{B},$$

as $\pi(f) = \oplus_{i=1}^m [f]_{p_i} \oplus_{j=1}^r [f]_{q_j}$, where $[f]_p$ is the residue class of f in the appropriate algebra. Then

$$\pi^* : \frac{\mathbb{R}[x]}{\ker \pi} \longrightarrow \mathcal{B}$$

is a monomorphism. Using Lemma 1.7 and (2.1) we get

$$\begin{aligned}
\ker \pi &= \mathbb{R}[x] \cap \bigcap_{i=1}^m (J_{\mathbb{R}} + m_{\mathbb{R}, p_i}^k) \cap \bigcap_{j=1}^r (J_{\mathbb{C}} + m_{\mathbb{C}, q_j}^k) = \\
&= \mathbb{R}[x] \cap \bigcap_{i=1}^m (J_{\mathbb{R}} + m_{\mathbb{R}, p_i}^k)_{\mathbb{C}} \cap \bigcap_{j=1}^r (J_{\mathbb{C}} + m_{\mathbb{C}, q_j}^k) = \\
&= \mathbb{R}[x] \cap \bigcap_{i=1}^m (J_{\mathbb{C}} + m_{\mathbb{C}, p_i}^k) \cap \bigcap_{j=1}^r (J_{\mathbb{C}} + m_{\mathbb{C}, q_j}^k) \cap \bigcap_{j=1}^r (J_{\mathbb{C}} + m_{\mathbb{C}, \bar{q}_j}^k) = \\
&= \mathbb{R}[x] \cap (J_{\mathbb{C}} : I_{\mathbb{C}}) = J_{\mathbb{R}} : I_{\mathbb{R}}.
\end{aligned}$$

By (2.2) we have

$$\begin{aligned}
\dim_{\mathbb{R}} \mathcal{B} &= \sum_{i=1}^m \dim_{\mathbb{R}} \mathcal{A}_{\mathbb{R}, p_i} + \sum_{j=1}^r \dim_{\mathbb{R}} \mathcal{A}_{\mathbb{C}, q_j} = \\
&= \sum_{i=1}^m \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C}, p_i} + 2 \cdot \sum_{j=1}^r \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C}, q_j} = \\
&= \sum_{i=1}^m \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C}, p_i} + \sum_{j=1}^r \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C}, q_j} + \sum_{j=1}^r \dim_{\mathbb{C}} \mathcal{A}_{\mathbb{C}, \bar{q}_j}.
\end{aligned}$$

By Proposition 1.8, it equals

$$\dim_{\mathbb{C}} \frac{\mathbb{C}[x]}{J_{\mathbb{C}} : I_{\mathbb{C}}} = \dim_{\mathbb{R}} \frac{\mathbb{R}[x]}{J_{\mathbb{R}} : I_{\mathbb{R}}}.$$

So $\dim_{\mathbb{R}} \mathcal{B} = \dim_{\mathbb{R}} \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}})$ and then $\pi^* : \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}}) \rightarrow \mathcal{B}$ is an isomorphism. $\square$

3 Bilinear forms

Let $\mathbb{K}$ denote either $\mathbb{R}$ or $\mathbb{C}$. For $p \in \mathbb{K}^n$, let $\mathcal{O}_{\mathbb{K}, p}$ denote the ring of germs at p of analytic functions $\mathbb{K}^n \rightarrow \mathbb{K}$. There is a natural homomorphism

$$\eta : \mathbb{K}[x] = \mathbb{K}[x_1, \dots, x_n] \rightarrow \mathcal{O}_{\mathbb{K}, p}.$$

Let $m_{\mathbb{K}, p} = \{f \in \mathbb{K}[x] : f(p) = 0\}$ be the maximal ideal in $\mathbb{K}[x]$ associated with p .

Let $S_{\mathbb{R}}$ be an ideal in $\mathbb{R}[x]$, let $S_{\mathbb{C}}$ denote the ideal in $\mathbb{C}[x]$ generated by $S_{\mathbb{R}}$, and let

$$V(S_{\mathbb{K}}) = \{p \in \mathbb{K}^n : f(p) = 0 \text{ for all } f \in S_{\mathbb{K}}\}.$$

Take $h_1, \dots, h_n \in S_{\mathbb{R}}$, the ideal $J_{\mathbb{K}}$ generated by $h_1, \dots, h_n$, and a polynomial mapping $H_{\mathbb{K}} = (h_1, \dots, h_n) : \mathbb{K}^n \rightarrow \mathbb{K}^n$. Then $J_{\mathbb{K}} \subset S_{\mathbb{K}}$ and $V(S_{\mathbb{K}}) \subset V(J_{\mathbb{K}}) = H_{\mathbb{K}}^{-1}(0)$. In particular, points isolated in $H_{\mathbb{C}}^{-1}(0)$ are isolated in $V(S_{\mathbb{K}})$.

Take $p \in V(S_{\mathbb{K}})$. Let $S_{\mathbb{K}, p}$ (resp. $J_{\mathbb{K}, p}$) denote the ideal in $\mathcal{O}_{\mathbb{K}, p}$ generated by $S_{\mathbb{K}}$ (resp. $J_{\mathbb{K}}$), and let $\mathcal{A}'_{\mathbb{K}, p} = \mathcal{O}_{\mathbb{K}, p}/S_{\mathbb{K}, p}$. Clearly $\mathcal{A}'_{\mathbb{K}, p}$ is a $\mathbb{K}$ -algebra and $\eta(S_{\mathbb{K}}) \subset S_{\mathbb{K}, p}$.

Assume that $S_{\mathbb{K}, p} = J_{\mathbb{K}, p}$ so that $\mathcal{A}'_{\mathbb{K}, p} = \mathcal{O}_{\mathbb{K}, p}/J_{\mathbb{K}, p}$. Then $\dim_{\mathbb{K}} \mathcal{A}'_{\mathbb{K}, p} < \infty$ if and only if p is isolated in $H_{\mathbb{C}}^{-1}(0)$. If that is the case then p is isolated in $V(S_{\mathbb{R}})$, and $\eta(m_{\mathbb{K}, p}^k) \subset S_{\mathbb{K}, p}$ for all k large enough.

Lemma 3.1 *If $p \in V(S_{\mathbb{K}})$ is isolated in $H_{\mathbb{C}}^{-1}(0)$ and $S_{\mathbb{K},p} = J_{\mathbb{K},p}$, then η induces an isomorphism of $\mathbb{K}$ -algebras*

$$\eta : \mathbb{K}[x]/(S_{\mathbb{K}} + m_{\mathbb{K},p}^k) \longrightarrow \mathcal{A}'_{\mathbb{K},p}$$

for all k large enough.

□

For p and k as above, put

$$\mathcal{A}_{\mathbb{K},p} = \mathbb{K}[x]/(S_{\mathbb{K}} + m_{\mathbb{K},p}^k),$$

so that $\mathcal{A}_{\mathbb{K},p}$ is isomorphic to $\mathcal{A}'_{\mathbb{K},p}$. In particular, $\mathcal{A}_{\mathbb{K},p}$ does not depend on k , if k is large enough, and $\dim_{\mathbb{K}} \mathcal{A}_{\mathbb{K},p} < \infty$.

Applying the formula for the local topological degree by Eisenbud and Levine [3] and Khimshiashvili [5, 6], and the theory of Frobenius algebras (see [1, 7, 9]) one may prove some properties, presented in [10], of bilinear forms on the algebra $\mathcal{O}_{\mathbb{K},p}/J_{\mathbb{K},p} = \mathcal{A}'_{\mathbb{K},p}$. As $\mathcal{A}'_{\mathbb{K},p}$ is isomorphic to $\mathcal{A}_{\mathbb{K},p}$, they hold true for the algebra $\mathcal{A}_{\mathbb{K},p}$.

Let points $p_1, \dots, p_w \in V(S_{\mathbb{C}})$ be as in Lemma 3.1. One may assume that $p_1, \dots, p_m \in \mathbb{R}^n$ and $p_{m+1}, \dots, p_w \in \mathbb{C}^n \setminus \mathbb{R}^n$. Denote

$$\mathcal{B} = \oplus_{i=1}^m \mathcal{A}_{\mathbb{R},p_i} \oplus_{j=m+1}^w \mathcal{A}_{\mathbb{C},p_j}.$$

Obviously, $\mathcal{B}$ is a finite dimensional $\mathbb{R}$ -algebra.

Let $u \in \mathbb{R}[x]$, and let $\varphi : \mathcal{B} \longrightarrow \mathbb{R}$ be a linear functional. Then there are bilinear symmetric forms $\Phi, \Psi : \mathcal{B} \times \mathcal{B} \longrightarrow \mathbb{R}$ given by $\Phi(a, b) = \varphi(ab)$ and $\Psi(a, b) = \varphi(uab)$.

One may define signature Φ , and similarly signature Ψ , as the dimension of a maximal subspace of $\mathcal{B}$ on which Φ is positive definite minus the dimension of that one on which Φ is negative definite.

Let $\det[\Phi]$ (resp. $\det[\Psi]$) denote the determinant of the matrix of Φ (resp. Ψ), with respect to some basis of $\mathcal{B}$. The sign of the determinant does not depend on the choice of a basis.

Theorem 3.2 [10, Theorem 2.3, p. 306] *Let points $p_1, \dots, p_w \in V(S_{\mathbb{C}}) \subset H_{\mathbb{C}}^{-1}(0)$ be as in Lemma 3.1. Suppose that $\det[\Psi] \neq 0$. Then $\det[\Phi] \neq 0$, $u(p_i) \neq 0$ for $1 \leq i \leq m$, and*

$$\sum \deg_p H_{\mathbb{R}} \equiv \dim_{\mathbb{R}} \mathcal{B} + 1 + (\operatorname{sgn} \det[\Phi] + \operatorname{sgn} \det[\Psi])/2 \pmod{2},$$

where $p \in \{p_1, \dots, p_m\} \cap \{u > 0\}$ and $\deg_p H_{\mathbb{R}}$ denotes the local topological degree of $H_{\mathbb{R}}$ at p .

□

For $x = (x_1, \dots, x_n)$, $x' = (x'_1, \dots, x'_n)$, and $1 \leq i, j \leq n$ define

$$T_{ij}(x, x') = \frac{h_i(x'_1, \dots, x'_{j-1}, x_j, \dots, x_n) - h_i(x'_1, \dots, x'_j, x_{j+1}, \dots, x_n)}{x_j - x'_j}.$$

It is easy to see that each T_{ij} extends to a polynomial, thus we may assume that

$$T_{ij} \in \mathbb{R}[x, x'] = \mathbb{R}[x_1, \dots, x_n, x'_1, \dots, x'_n].$$

There is the natural projection $\mathbb{R}[x, x'] \longrightarrow \mathcal{B} \otimes \mathcal{B}$ given by

$$x_1^{\alpha_1} \cdots x_n^{\alpha_n} (x'_1)^{\beta_1} \cdots (x'_n)^{\beta_n} \mapsto x_1^{\alpha_1} \cdots x_n^{\alpha_n} \otimes (x'_1)^{\beta_1} \cdots (x'_n)^{\beta_n}.$$

Let T denote the image of $\det[T_{ij}(x, x')]$ in $\mathcal{B} \otimes \mathcal{B}$.

Put $d = \dim_{\mathbb{R}} \mathcal{B}$. Assume that $e_1, \dots, e_d$ form a basis in $\mathcal{B}$. So $\dim_{\mathbb{R}} \mathcal{B} \otimes \mathcal{B} = d^2$ and $e_i \otimes e_j$, for $1 \leq i, j \leq d$, form a basis in $\mathcal{B} \otimes \mathcal{B}$. Hence there are $t_{ij} \in \mathbb{R}$ such that

$$T = \sum_{i,j=1}^d t_{ij} e_i \otimes e_j = \sum_{i=1}^d e_i \otimes \hat{e}_i,$$

where $\hat{e}_i = \sum_{j=1}^d t_{ij} e_j$. Elements $\hat{e}_1, \dots, \hat{e}_d$ form a basis in $\mathcal{B}$. So there are $A_1, \dots, A_d \in \mathbb{R}$ such that

$$1 = A_1 \hat{e}_1 + \dots + A_d \hat{e}_d \text{ in } \mathcal{B}.$$

Definition. For $f = a_1 e_1 + \dots + a_d e_d \in \mathcal{B}$ define $\varphi_T(f) = a_1 A_1 + \dots + a_d A_d$. Hence $\varphi_T : \mathcal{B} \longrightarrow \mathbb{R}$ is a linear functional.

Let Φ_T be the bilinear form on $\mathcal{B}$ given by $\Phi_T(a, b) = \varphi_T(ab)$.

Theorem 3.3 [10, Theorem 1.5, p. 304] *The form Φ_T is non-degenerate and*

$$\sum_{i=1}^m \deg_{p_i} H_{\mathbb{R}} = \text{signature } \Phi_T.$$

□

4 Topological degree

Let $h_1, \dots, h_n \in \mathbb{R}[x_1, \dots, x_n]$, and let $H_{\mathbb{K}} = (h_1, \dots, h_n) : \mathbb{K}^n \longrightarrow \mathbb{K}^n$. Denote by $J_{\mathbb{K}}$ the ideal in $\mathbb{K}[x]$ generated by $h_1, \dots, h_n$, so that $H_{\mathbb{C}}^{-1}(0) = V(J_{\mathbb{C}})$.

Assume that there is an ideal $I_{\mathbb{R}}$ such that $J_{\mathbb{C}} \subset I_{\mathbb{C}}$, $\dim_{\mathbb{R}} \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}}) < \infty$, and $(J_{\mathbb{R}} : I_{\mathbb{R}}) + I_{\mathbb{R}} = \mathbb{R}[x]$.

Put $S_{\mathbb{R}} = J_{\mathbb{R}} : I_{\mathbb{R}}$. Hence $V(S_{\mathbb{C}})$ is finite, and by Lemma 1.4

$$H_{\mathbb{C}}^{-1}(0) \setminus V(I_{\mathbb{C}}) = V(J_{\mathbb{C}}) \setminus V(I_{\mathbb{C}}) = V(S_{\mathbb{C}}).$$

Hence each $p \in V(J_{\mathbb{C}}) \setminus V(I_{\mathbb{C}})$ is isolated in $H_{\mathbb{C}}^{-1}(0)$. By Corollary 1.7, if k is large enough then the ideal $S_{\mathbb{K},p} \subset \mathcal{O}_{\mathbb{K},p}$ generated by $S_{\mathbb{R}}$ equals the ideal generated by $J_{\mathbb{K}} + m_{\mathbb{K},p}^k$.

Let $J_{\mathbb{K},p}$ denote the ideal in $\mathcal{O}_{\mathbb{K},p}$ generated by $J_{\mathbb{R}}$. Since $J_{\mathbb{K},p}$ has an algebraically isolated zero at p , the local Nullstellensatz implies that $m_{\mathbb{K},p}^k \subset J_{\mathbb{K},p}$, so that $S_{\mathbb{K},p} = J_{\mathbb{K},p} + m_{\mathbb{K},p}^k = J_{\mathbb{K},p}$. Hence each point $p \in V(S_{\mathbb{C}}) = H_{\mathbb{C}}^{-1}(0) \setminus V(I_{\mathbb{C}})$ satisfies the assumptions of Lemma 3.1.

Put $H_{\mathbb{R}}^{-1}(0) \setminus V(I_{\mathbb{R}}) = \{p_1, \dots, p_m\}$ and

$$(H_{\mathbb{C}}^{-1}(0) \setminus V(I_{\mathbb{C}})) \setminus \mathbb{R}^n = \{q_1, \overline{q_1}, \dots, q_r, \overline{q_r}\}.$$

By Theorem 2.3, $\mathcal{A} := \mathbb{R}[x]/S = \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}})$ and

$$\mathcal{B} = \oplus_{i=1}^m \mathcal{A}_{\mathbb{R}, p_i} \oplus_{j=1}^r \mathcal{A}_{\mathbb{C}, q_j}$$

are isomorphic.

As a consequence of Theorem 3.2 we get

Theorem 4.1 *Assume that $\dim_{\mathbb{R}} \mathcal{A} < \infty$, $u \in \mathbb{R}[x]$ and $\varphi : \mathcal{A} \rightarrow \mathbb{R}$ is a linear functional. Let bilinear symmetric forms $\Phi, \Psi : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{R}$ be given by $\Phi(a, b) = \varphi(ab)$ and $\Psi(a, b) = \varphi(uab)$.*

Suppose that $\det[\Psi] \neq 0$. Then $\det[\Phi] \neq 0$, $u(p) \neq 0$ for each $p \in H_{\mathbb{R}}^{-1}(0) \setminus V(I_{\mathbb{R}})$, and

$$\sum \deg_p H_{\mathbb{R}} \equiv \dim_{\mathbb{R}} \mathcal{A} + 1 + (\operatorname{sgn} \det[\Phi] + \operatorname{sgn} \det[\Psi])/2 \pmod{2},$$

where $p \in H_{\mathbb{R}}^{-1}(0) \cap \{u > 0\} \setminus V(I_{\mathbb{R}})$.

□

The same way as in Section 3 one may construct the bilinear symmetric form Φ_T on $\mathcal{A} = \mathbb{R}[x]/(J_{\mathbb{R}} : I_{\mathbb{R}})$. As a consequence of Theorem 3.3 we get

Theorem 4.2 *The form Φ_T is non-degenerate and*

$$\operatorname{signature} \Phi_T = \sum \deg_p H_{\mathbb{R}},$$

where $p \in H_{\mathbb{R}}^{-1}(0) \setminus V(I_{\mathbb{R}})$.

□

Let Ψ_T be the bilinear form on $\mathcal{A}$ given by $\Psi_T(a, b) = \varphi_T(uab)$. Using the same arguments as in [10, Theorem 1.5, p. 304] one may prove

Theorem 4.3 *Ψ_T is non-degenerate if and only if $u(p) \neq 0$ for each $p \in H_{\mathbb{C}}^{-1}(0) \setminus V(I_{\mathbb{C}})$. If that is the case then*

$$\sum \deg_p H_{\mathbb{R}} = \frac{1}{2}(\operatorname{signature} \Phi_T + \operatorname{signature} \Psi_T),$$

where $p \in H_{\mathbb{R}}^{-1}(0) \cap \{u > 0\} \setminus V(I_{\mathbb{R}})$.

5 Immersions

Let M be an m -dimensional manifold. A C^1 map $g : M \rightarrow \mathbb{R}^k$ is called an immersion, if for each $p \in M$ the rank of $Dg(p)$ equals m .

Assume that m is even, $k = 2m$ and M is compact and oriented. Define

$$G : M \times M \rightarrow \mathbb{R}^{2m} \quad \text{as} \quad G(x, y) = g(x) - g(y).$$

Set $\Delta = \{(p, p) : p \in M\} \subset M \times M$. Since g is an immersion, Δ is isolated in $G^{-1}(0)$, and so $G^{-1}(0) \setminus \Delta$ is a compact subset of $M \times M \setminus \Delta$. There exists $(N, \partial N)$ — a compact $2m$ -dimensional oriented manifold with boundary, such that

$$N \subset M \times M \setminus \Delta \quad \text{and} \quad G^{-1}(0) \setminus \Delta \subset N \setminus \partial N.$$

Denote by $d(G)$ the topological degree of the mapping

$$\partial N \ni (x, y) \mapsto \frac{G(x, y)}{\|G(x, y)\|} \in S^{2m-1}.$$

Of course, $d(G)$ does not depend on the choice of N . In particular, if $G^{-1}(0) \setminus \Delta$ is finite then

$$d(G) = \sum \deg_z G, \quad \text{where } z \in G^{-1}(0) \setminus \Delta,$$

and $\deg_z G$ denotes the local topological degree of G at z .

Whitney has introduced in [12] an intersection number $I(g)$ for the immersion g . According to the Lashof and Smale [8, Theorem 3.1],

$$d(G) = 2I(g).$$

Now assume that $f = (f_1, \dots, f_n) : \mathbb{R}^{n+m} \longrightarrow \mathbb{R}^n$ is a C^1 mapping, such that $M = f^{-1}(0)$ and M is a complete intersection, i.e. for each $p \in M$ the rank of $Df(p)$ equals n .

We shall say that vectors $v_1, \dots, v_m \in T_p M$ are well oriented if vectors $\nabla f_1(p), \dots, \nabla f_n(p), v_1, \dots, v_m$ are well oriented in $\mathbb{R}^{n+m}$. Put

$$F(x, y) = (f(x), f(y)) : \mathbb{R}^{n+m} \times \mathbb{R}^{n+m} \longrightarrow \mathbb{R}^n \times \mathbb{R}^n.$$

Then $F^{-1}(0) = M \times M$ is a complete intersection. The orientation of $F^{-1}(0)$ defined the way as above is the same as the orientation of the product $M \times M$.

Let

$$\bar{g} = (g_1, \dots, g_{2m}) : \mathbb{R}^{n+m} \longrightarrow \mathbb{R}^{2m}$$

be a C^1 mapping. Put $g = \bar{g}|_M$. It easy to verify that

$$\text{rank} \begin{bmatrix} Dg(p) \end{bmatrix} = \text{rank} \begin{bmatrix} D\bar{g}(p) \\ Df(p) \end{bmatrix} - n$$

at each point $p \in M$, so we have

Lemma 5.1 $g = \bar{g}|_M : M \longrightarrow \mathbb{R}^{2m}$ is an immersion if and only if at each point $p \in M$

$$\text{rank} \begin{bmatrix} D\bar{g}(p) \\ Df(p) \end{bmatrix} = n + m.$$

□

Example 5.2 Let $f = x^2 + y^2 + z^2 - r^2$, and

$$\bar{g} = (x, y, xz, yz) : \mathbb{R}^3 \longrightarrow \mathbb{R}^4,$$

i.e. $m = 2, n = 1$. Then $M = f^{-1}(0)$ is the 2-dimensional sphere $S^2(r)$ of radius r . As

$$\text{rank} \begin{bmatrix} D\bar{g} \\ Df \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ z & 0 & x \\ 0 & z & y \\ 2x & 2y & 2z \end{bmatrix}$$

has a non-zero (3×3) -minor at each point $p \in \mathbb{R}^3 \setminus \{0\}$, then $g = \bar{g}|_{S^2(r)}$ is an immersion for every $r > 0$.

Let

$$\overline{G}(x, y) = \overline{g}(x) - \overline{g}(y).$$

Then $G(x, y) = g(x) - g(y) = \overline{G}|_{M \times M} = \overline{G}|_{F^{-1}(0)}$. Put

$$\begin{aligned} H(x, y) &= (F(x, y), \overline{G}(x, y)) = \\ &= (f_1(x), \dots, f_n(x), f_1(y), \dots, f_n(y), g_1(x) - g_1(y), \dots, g_{2m}(x) - g_{2m}(y)). \end{aligned}$$

Then $H : \mathbb{R}^{n+m} \times \mathbb{R}^{n+m} \longrightarrow \mathbb{R}^{2n+2m}$, and $(p, q) \in H^{-1}(0)$ if and only if $(p, q) \in M \times M$ and $G(p, q) = 0$. By [10, Lemma 3.2], $z = (p, q) \in M \times M$ is isolated in $G^{-1}(0)$ if and only if z is isolated in $H^{-1}(0)$, and if that is the case then

$$\deg_z G = \deg_z H.$$

Let $\Delta = \{(p, p) | p \in \mathbb{R}^{n+m}\}$ be the diagonal in $\mathbb{R}^{n+m} \times \mathbb{R}^{n+m}$. Then

$$z = (p, q) \in H^{-1}(0) \setminus \Delta \quad \text{if and only if} \quad z = (p, q) \in G^{-1}(0) \setminus \Delta.$$

So we get

Proposition 5.3 *Suppose that m is even, and*

- (a) $M = f^{-1}(0)$ is an oriented compact m -dimensional complete intersection,
- (b) $g = \overline{g}|_M : M \longrightarrow \mathbb{R}^{2m}$ is an immersion,
- (c) $H^{-1}(0) \setminus \Delta$ is finite.

Then $2I(g) = d(G) = \sum \deg_z H$, where $z \in H^{-1}(0) \setminus \Delta$.

□

A homotopy $h_t : M \longrightarrow \mathbb{R}^{2m}$ is called a *regular homotopy*, if at each stage it is an immersion and the induced homotopy of the tangent bundle is continuous.

As in [12] we say that an immersion $g : M \longrightarrow \mathbb{R}^{2m}$ has a *regular self-intersection* at the point $g(p) = g(q)$ if

$$Dg(p)T_p M + Dg(q)T_q M = \mathbb{R}^{2m}.$$

That is so if and only if $\det[DH(p, q)] \neq 0$.

If g has only regular self-intersections, and there are no triple points $g(p) = g(q) = g(w)$, then we say that g is completely regular. If m is odd, then one can define the *intersection number* of a completely regular immersion as the number of its self-intersections modulo 2.

By [12, Theorem 2], if M is compact then the intersection number is invariant under regular homotopies. As in [11, §8], for any immersion $g : M \longrightarrow \mathbb{R}^{2m}$ there exists a completely regular immersion $\tilde{g} : M \longrightarrow \mathbb{R}^{2m}$ which is arbitrarily close to g in C^1 -topology and there is a regular homotopy between g and $\tilde{g}$. Thus if $m > 1$ is odd then one can define the intersection number $I(g)$ as the number of self-intersections of $\tilde{g}$ modulo 2.

Suppose that g has a finite number of self-intersections, i.e. $H^{-1}(0) \setminus \Delta$ is finite. The immersion has a self-intersection $g(p) = g(q)$ if and only if (p, q) and (q, p) belong to $H^{-1}(0) \setminus \Delta$. There exists a linear function

$$u(x, y) = a_1(x_1 - y_1) + \dots + a_{n+m}(x_{n+m} - y_{n+m})$$

which does not vanish at any point in $H^{-1}(0) \setminus \Delta$, so that $u(p, q) > 0$ if and only if $u(q, p) < 0$. Then each self-intersection is represented by a single point in $H^{-1}(0) \cap \{u > 0\} \setminus \Delta$.

Proposition 5.4 *Suppose that $m > 1$ is odd, and*

- (a) $M = f^{-1}(0)$ is an oriented, compact m -dimensional complete intersection,
- (b) $g = \bar{g}|M: M \longrightarrow \mathbb{R}^{2m}$ is an immersion,
- (c) $H^{-1}(0) \setminus \Delta$ is finite,
- (d) $u(x, y) = a_1(x_1 - y_1) + \dots + a_{n+m}(x_{n+m} - y_{n+m})$ is such that $u(p, q) \neq 0$ for each $(p, q) \in H^{-1}(0) \setminus \Delta$.

Then

$$I(g) \equiv \sum \deg_z H \pmod{2},$$

where $z \in H^{-1}(0) \cap \{u > 0\} \setminus \Delta$.

Proof. Put

$$\begin{aligned} \tilde{H}(x, y) &= \\ &= (f_1(x), \dots, f_n(x), f_1(y), \dots, f_n(y), \tilde{g}_1(x) - \tilde{g}_1(y), \dots, \tilde{g}_{2m}(x) - \tilde{g}_{2m}(y)). \end{aligned}$$

Let $B_z \subset \mathbb{R}^{n+m} \times \mathbb{R}^{n+m}$ denote a small ball centered at $z \in H^{-1}(0) \cap \{u > 0\} \setminus \Delta$. If $\tilde{g}$ is close enough to g , then self-intersections of $\tilde{g}$ are represented by points in the set $\tilde{H}^{-1}(0) \cap \{u > 0\} \setminus \Delta$, which is a subset of the union of all B_z . As self-intersections of $\tilde{g}$ are regular,

$$I(\tilde{g}) \equiv \sum_w \operatorname{sgn} \det[D\tilde{H}(w)] \pmod{2},$$

where $w \in \tilde{H}^{-1}(0) \cap \{u > 0\} \setminus \Delta$. Each point w belongs to some B_z , hence

$$I(g) = I(\tilde{g}) \equiv \sum_z \sum_w \operatorname{sgn} \det[D\tilde{H}(w)] \pmod{2},$$

where $z \in H^{-1}(0) \cap \{u > 0\} \setminus \Delta$ and $w \in \tilde{H}^{-1}(0) \cap B_z$. Since $\tilde{H}$ is close to H in a neighbourhood of $M \times M$,

$$I(g) \equiv \sum_z \deg_z H \pmod{2}.$$

□

If $f_1, \dots, f_n, g_1, \dots, g_{2m}$ are polynomials then $H = (h_1, \dots, h_{2n+2m})$ is a polynomial mapping. Let $J_{\mathbb{R}}$ denote the ideal in $\mathbb{R}[x, y] = \mathbb{R}[x_1, \dots, x_{n+m}, y_1, \dots, y_{n+m}]$ generated by $h_1, \dots, h_{2n+2m}$, and $I_{\mathbb{R}}$ the one generated by

$$f_1(x), \dots, f_n(x), f_1(y), \dots, f_n(y), x_1 - y_1, \dots, x_{n+m} - y_{n+m}.$$

It is easy to verify that $J_{\mathbb{R}} \subset I_{\mathbb{R}}$. Then

$$V(J_{\mathbb{R}}) = H^{-1}(0), \quad V(I_{\mathbb{R}}) = M \times M \cap \Delta.$$

Let $Q = \mathbb{R}[x, y]/J_{\mathbb{R}}$. If $m \geq 1$ and $M \neq \emptyset$, then $M \times M \cap \Delta \subset H^{-1}(0)$ is infinite, so $\dim_{\mathbb{R}} Q = \infty$ and we cannot apply methods presented in [10] so as to compute $\sum_p \deg_p H$.

Let $\mathcal{A} = \mathbb{R}[x, y]/(J_{\mathbb{R}} : I_{\mathbb{R}})$. Suppose that $\dim_{\mathbb{R}} \mathcal{A} < \infty$ and $(J_{\mathbb{R}} : I_{\mathbb{R}}) + I_{\mathbb{R}} = \mathbb{R}[x, y]$. Then $H^{-1}(0) \setminus \Delta$ is finite, so that the immersion g has a finite set of self-intersections.

Let $\Phi_T : \mathcal{A} \times \mathcal{A} \longrightarrow \mathbb{R}$ be the bilinear form constructed as in Section 3. As a consequence of Theorem 4.2 and Proposition 5.3 we get

Theorem 5.5 *If m is even, then*

$$I(g) = \frac{1}{2} \text{signature } \Phi_T.$$

□

Take any polynomial $u(x, y) = a_1(x_1 - y_1) + \dots + a_{n+m}(x_{n+m} - y_{n-m})$ and any linear form $\varphi : \mathcal{A} \longrightarrow \mathbb{R}$. Let $\Phi, \Psi : \mathcal{A} \times \mathcal{A} \longrightarrow \mathbb{R}$ be bilinear forms given by $\Phi(a, b) = \varphi(ab)$, $\Psi(a, b) = \varphi(uab)$. As a consequence of Theorem 4.1 and Proposition 5.4 we get

Theorem 5.6 *If $m > 1$ is odd and $\det[\Psi] \neq 0$, then*

$$I(g) \equiv \dim_{\mathbb{R}} \mathcal{A} + 1 + (\text{sgn } \det[\Phi] + \text{sgn } \det[\Psi])/2 \pmod{2}.$$

□

Example 5.7 *Let us consider the mapping*

$$g = (g_1, g_2, g_3, g_4) = (x_1, x_2, x_1x_3, x_2x_3) : \mathbb{R}^3 \longrightarrow \mathbb{R}^4.$$

As in Example 5.2, $g|_{S^2(1)}$ is an immersion. Put $f = x_1^2 + x_2^2 + x_3^2 - 1$. With the immersion $g|_{S^2(1)}$ we may associate the polynomial mapping

$$H = (h_1, \dots, h_6) : \mathbb{R}^3 \times \mathbb{R}^3 \longrightarrow \mathbb{R}^6$$

given by $h_1 = f(x_1, x_2, x_3)$, $h_2 = f(y_1, y_2, y_3)$ and $h_i = g_{i-2}(x_1, x_2, x_3) - g_{i-2}(y_1, y_2, y_3)$, for $i = 3, 4, 5, 6$.

Let $I_{\mathbb{R}}$ be the ideal in $\mathbb{R}[x_1, x_2, x_3, y_1, y_2, y_3]$ generated by $f(x), f(y), x_1 - y_1, x_2 - y_2, x_3 - y_3$ and let $J_{\mathbb{R}}$ be generated by $h_1, \dots, h_6$. Using SINGULAR — a computer algebra system for polynomial computations — one may check that $J_{\mathbb{R}} : I_{\mathbb{R}}$ is generated by $x_1, x_2, y_1, y_2, x_3 + y_3, y_3^2 - 1$, and then monomials $e_1 = 1$ and $e_2 = y_3$ form a basis of $\mathcal{A} = \mathbb{R}[x_1, \dots, y_3]/(J_{\mathbb{R}} : I_{\mathbb{R}})$. Then $\dim_{\mathbb{R}} \mathcal{A} = 2 < \infty$ and $(J_{\mathbb{R}} : I_{\mathbb{R}}) + I_{\mathbb{R}} = \mathbb{R}[x_1, x_2, x_3, y_1, y_2, y_3]$. One may check that

$$T = -8y_3y'_3 - 8 \quad \text{in } \mathcal{A} \otimes \mathcal{A}.$$

So $\hat{e}_1 = -8$ and $\hat{e}_2 = -8y'_3$. Clearly $1 = (-\frac{1}{8})\hat{e}_1 + 0\hat{e}_2$ in $\mathcal{A}$. Then $A_1 = -\frac{1}{8}$, $A_2 = 0$, so for any $a = a_1 + a_2y_3 = a_1e_1 + a_2e_2$ in $\mathcal{A}$, φ_T is given by $\varphi_T(a) = -\frac{a_1}{8}$. Then the matrix of Φ_T is given by

$$\begin{bmatrix} -\frac{1}{8} & 0 \\ 0 & -\frac{1}{8} \end{bmatrix},$$

so $\text{signature } \Phi_T = -2$, and as a consequence of Theorem 5.5 we get $I(g|_{S^2(1)}) = -1$.

Example 5.8 Let $g = (g_1, \dots, g_6) = (x_1, x_2, x_1x_3, x_2x_3, x_4, x_3x_4) : \mathbb{R}^4 \longrightarrow \mathbb{R}^6$. Using Lemma 5.1 it is easy to check that $g|_{S^3(1)}$ is an immersion. Put $f = x_1^2 + x_2^2 + x_3^2 + x_4^2 - 1$ and

$$\begin{aligned} H(x_1, \dots, y_4) &= (h_1, \dots, h_8) = \\ &= (f(x_1, \dots, x_4), f(y_1, \dots, y_4), g(x_1, \dots, x_4) - g(y_1, \dots, y_4)). \end{aligned}$$

Let $I_{\mathbb{R}} \subset \mathbb{R}[x_1, \dots, y_4]$ be the ideal generated by $f(x), f(y), x_1 - y_1, x_2 - y_2, x_3 - y_3, x_4 - y_4$, and $J_{\mathbb{R}}$ generated by $h_1, \dots, h_8$. One may check, using SINGULAR, that $x_1, x_2, x_4, y_1, y_2, y_4, x_3 + y_3, y_3^2 - 1$ generate $J_{\mathbb{R}} : I_{\mathbb{R}}$, and that monomials $e_1 = 1, e_2 = y_3$ form the basis of $\mathcal{A} = \mathbb{R}[x_1, \dots, y_4]/(J_{\mathbb{R}} : I_{\mathbb{R}})$. Then $\dim_{\mathbb{R}} \mathcal{A} = 2 < \infty$ and $(J_{\mathbb{R}} : I_{\mathbb{R}}) + I_{\mathbb{R}} = \mathbb{R}[x_1, \dots, y_4]$. Put $\varphi : \mathcal{A} \longrightarrow \mathbb{R}$ as

$$\varphi(a) = \varphi(a_1e_1 + a_2e_2) = a_2.$$

Take $u = x_3 - y_3$. Matrices of Φ and Ψ are given by

$$\begin{bmatrix} 0 & +1 \\ +1 & 0 \end{bmatrix}, \quad \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix}.$$

Of course $\det[\Psi] \neq 0$, and as consequence of Theorem 5.6 we get $I(g|_{S^3(1)}) \equiv 2 + 1 + \frac{1}{2}(-1 + 1) \equiv 1 \pmod{2}$.

Using similar methods we have computed some more difficult examples:

Example 5.9 $h(x_1, x_2, x_3) =$

$$= (2x_1x_2 + x_2, 2x_1x_3 + 4x_3, 4x_3^2 + 5x_2, 5x_2^2 + 4x_3)$$

is an immersion on the 2-dimensional sphere of radius $r = 10$. In that case $\dim_{\mathbb{R}} \mathcal{A} = 16$, and $I(h|_{S^2(10)}) = 0$.

Example 5.10 $h(x_1, x_2, x_3) =$

$$= (5x_2x_3 + x_3^2 + 3x_1, 4x_1^2 + 3x_3^2 + x_3, 2x_2^2 + 3x_2x_3 + 2x_1, x_2x_3 + 4x_3^2 + 3x_2)$$

is an immersion on the 2-dimensional spheres of radius $r = 1, 10$. In both cases $\dim_{\mathbb{R}} \mathcal{A} = 6$, and $I(h|_{S^2(1)}) = 0, I(h|_{S^2(10)}) = 1$.

Example 5.11 $h(x_1, x_2, x_3) =$

$$\begin{aligned} &= (3x_1x_2 + 2x_2^2 + 2x_1x_3 + 3x_1 + 5x_3, 2x_1x_2 + 5x_2^2 + 3x_2x_3 + x_1 + 2x_2, \\ &4x_1^2 + 4x_1x_3 + 5x_2x_3 + 3x_1 + 3x_3, 4x_2^2 + 3x_1x_3 + 4x_2x_3 + 4x_1 + 4x_3) \end{aligned}$$

is an immersion on the 2-dimensional spheres of radius $r = 1$. In that case $\dim_{\mathbb{R}} \mathcal{A} = 20$, and $I(h|_{S^2(1)}) = 1$.

Example 5.12 $h(x_1, x_2, x_3, x_4, x_5) =$

$$= (x_1x_2 + x_2, 3x_3x_5 + 2x_1, x_1^2 + x_2, x_3^2 + 3x_3, 3x_1x_5 + x_1, 4x_2x_5 + x_1, 2x_4^2 + x_4, x_3^2 + x_5)$$

is an immersion on the 4-dimensional spheres of radius $r = 1, 10$. In both cases $\dim_{\mathbb{R}} \mathcal{A} = 10$, and $I(h|_{S^4(1)}) = 0, I(h|_{S^4(10)}) = -1$.

Example 5.13 $h(x_1, x_2, x_3, x_4) =$

$$= (x_2x_4 + x_4, 2x_1x_4 + x_3, 3x_2x_4 + 4x_1, 3x_3x_4 + x_3, x_1x_2 + x_3, 2x_2x_3 + x_1)$$

is an immersion on the 3-dimensional sphere of radius $r = 1$. Take $u = 3(x_1 - y_1) + 5(x_2 - y_2) - 2(x_4 - y_4)$. In that case $\dim_{\mathbb{R}} \mathcal{A} = 18$, and $I(h|_{S^3(1)}) \equiv 1 \pmod{2}$.

References

- [1] E. Becker, J.P. Cardinal, M.F. Roy, and Z. Szafraniec, Multivariate Bezutians, Kronecker symbol and Eisenbud-Levine formula, Progress in Mathematics, vol.143, 1996, p. 79-104,
- [2] D. Cox, J. Little, D. O'Shea, Ideals, Varieties and Algorithms, Second Edition (1996),
- [3] D. Eisenbud, H.I. Levine, An algebraic formula for the degree of C^∞ map germ, Annals of Mathematics, vol. 106, 1977, p. 19-44,
- [4] G.-M. Greuel, G. Pfister, and H. Schönemann, SINGULAR 3.0.2. A Computer Algebra System for Polynomial Computations. Centre for Computer Algebra, University of Kaiserslautern (2006). <http://www.singular.uni-kl.de>,
- [5] G.M. Khimshiashvili, On the local degree of a smooth map, Soobshch. Akad. Nauk Gruz. SSR 85, 1977, p. 309-311,
- [6] G.M. Khimshiashvili, On the local degree of a smooth map, Trudy Tbilisi Math. Inst. 64, 1980, p. 105-124,
- [7] E. Kunz, Kähler Differentials, Vieweg Advanced Lectures in Mathematics, Braunschweig, Wiesbaden, 1986,
- [8] R. Lashof and S. Smale, On the immersion of manifolds in euclidean spaces, Annals of Mathematics, vol.68, no.3, 1958, p. 562-583,
- [9] G. Scheja and U. Storch, Über Spurfunktionen bei vollständigen Durchschnitten, J. Reine Angew Math. 278/279, 1975, p. 174-190,
- [10] Z. Szafraniec, Topological degree and quadratic forms, Journal of Pure and Applied Algebra 141, 1999, p. 299-314,
- [11] H. Whitney, Differentiable manifolds, Annals of Mathematics, vol. 37, no. 3, 1936, p. 645-680,
- [12] H. Whitney, The self-intersections of a smooth n -manifold in $2n$ -space, Annals of Mathematics vol. 45, no. 2, 1944, p. 220-246.